

Main characteristics of Westerwolds ryegrass varieties¹

| Ras | Year of admission | Average date of flowering | Juvenile growth (1-9) ² | Regrowth (1-9) ² | Rust-resistance (1-9) ² | DM-yield as main crop ³ | DM-yield as second crop ³ |
|-----------------------------|-------------------|---------------------------|------------------------------------|-----------------------------|------------------------------------|------------------------------------|--------------------------------------|
| <i>Varieties diploid</i> | | | | | | | |
| MENDOZA | 2001 | late / very late | 7,8 | 7,8 | 6,5 | 99 | 95 |
| MELSEMPER | 2011 | semi late | 7,6 | 8,6 | 7,5 | 100 | 103 |
| MELWORLD | 2004 | semi late | 7,5 | 8,6 | 7,2 | 105 | 101 |
| <i>Varieties tetraploid</i> | | | | | | | |
| LEMNOS | 1992 | semi late | 8,4 | 7,3 | 6,4 | 96 | 101 |
| MELMONDO ⁴ | 2003 | late / very late | 8,2 | - | 5,8 | - | 100 |

¹ Reproduction of the complete table from the descriptive variety list provided that the source is acknowledged, counterfeiting prohibited.

² The higher the number, the better

³ 100 = average of all varieties on the Belgian variety list

⁴ Only tested as second crop